

Research Article

Assessing Radiologic Findings of Chest Radiographs in Patients with Trauma Admitted to the Emergency Room

Mahsa Mohammadian¹, Alireza Shakibafard², Mahsa Akhavan¹, Zahra Ghahramani^{3*}, Shahram Paydar³

¹General Practitioner, Shiraz University of Medical Sciences, Shiraz, Iran

²Department of Radiology, Shiraz University of Medical Sciences, Shiraz, Iran

³Trauma Research Center, Shahid Rajaei (Emtiaz) Trauma Hospital, Shiraz University of Medical Sciences, Shiraz, Iran

*Corresponding author: Zahra Ghahramani, Trauma Research Center, Shahid Rajaei Trauma Hospital, Shiraz University of Medical Sciences, Shiraz, Iran. Tel: +987136360697; Fax: +987136254206; Email: ghahreman2@yahoo.co.nz.

Citation: Mohammadian M, Shakibafard A, Akhavan M, Ghahramani Z, Paydar S (2017) Assessing Radiologic Findings of Chest Radiographs in Patients with Trauma Admitted to the Emergency Room. Emerg Med Inves: EMIG-155.

Received Date: 5 August, 2017; **Accepted Date:** 22 September, 2017; **Published Date:** 29 September, 2017

Abstract

Background: According to Advanced Trauma Life Support, chest radiography must be performed during the initial evaluation of patients with trauma. We studied the CXR performed in the emergency room of Rajaie Hospital to determine its usefulness.

Methods: In this prospective study, patients who referred with high-energy trauma from December 2013 until April 2014 were recruited. Their demographic characteristics, including age, gender, and cause of trauma were recorded. Meticulous medical history was taken from all patients and they were examined by emergency medicine specialist. Simple radiographic CXR was performed for selective patients and image findings and their mediastinal width were recorded. All statistical analysis was performed using SPSS software version 20.0 and P-value less than 0.05 was considered significant.

Results: Of the total 790 patients assessed, 137 patients were female (17.3%) and 655 were male (82.7%). Mean age of patients was 35.13±17.01 (ranging from 12 to 91); Mean mediastinal width was 80.9±11.45 mm (range: 49.2-142.29). The most common causes of trauma included car-to-patient accident in 131 patients (16.5%), motor-to-car accident in 128 patients (16.2%), car turn-over in 103 (13%), falling down in 93 (11.7%), and stab wound 88 (11.1%). The most common pathologic finding included rib fractures (42.7%), pneumothorax (11.6%), abnormal diaphragmatic findings (10.3%), and hemothorax (8.3%).

Conclusion: The results of the current study, in accordance to previous studies, suggest rib fractures, pneumothorax, abnormal diaphragmatic findings, and hemothorax as the most frequent findings in CXR that need to be assessed meticulously. Moreover, the majority of patients were young males and the most frequent causes of trauma included car accidents, falling down, and stab wound. Therefore, paying attention to their diagnosis and treatment may increase the survival of this important group of patients.

Keywords: Chest X-Ray; Multiple Trauma; Radiologic Findings

Introduction

Multiple trauma is an important cause of death worldwide [1], which is also a leading cause of mortality and morbidity in Iran after cardiovascular injuries [2]. Although it is estimated that a number of cases pass away before reaching medical centers, it is essential to diagnose and manage the patients referring to trauma

centers rapidly and properly to increase their survival rate [3]. Among various traumas the multiple traumas that a patient might experience, chest trauma is ranked third and requires careful assessment [4].

Performing Chest X-Ray (CXR) in patients with multiple trauma is recommended by Advanced Trauma Life Support (ATLS) since 1992 [5] and is performed in many countries as the standard trauma care [6]. It is also an easy and fast imaging technique with low radiation dose [7] and can be interpreted rapidly. Studies have

Citation: Mohammadian M, Shakibafard A, Akhavan M, Ghahramani Z, Paydar S (2017) Assessing Radiologic Findings of Chest Radiographs in Patients with Trauma Admitted to the Emergency Room. *Emerg Med Inves: EMIG-155*.

RIGHT	No.	%	LEFT	No.	%	Total percentage
Pneumothorax	38	4.8	Pneumothorax	54	6.8	11.6
Hemothorax	65	8.2	Hemothorax	1	0.1	8.3
Diaphragm Abnormalities	25	3.2	Diaphragm Abnormalities	56	7.1	10.3
Clavicle Fx	28	3.5	Clavicle Fx	25	3.2	6.7
1 st Rib Fx	5	0.6	1 st Rib Fx	4	0.5	1.1
2 nd Rib Fx	16	2.0	2 nd Rib Fx	13	1.6	3.6
3 rd Rib Fx	21	2.7	3 rd Rib Fx	24	2.7	5.4
4 th Rib Fx	24	3.0	4 th Rib Fx	24	3.0	6.0
5 th Rib Fx	27	3.4	5 th Rib Fx	20	2.5	6.9
6 th Rib Fx	25	3.2	6 th Rib Fx	20	2.5	5.7
7 th Rib Fx	19	2.4	7 th Rib Fx	22	2.8	5.2
8 th Rib Fx	15	1.9	8 th Rib Fx	18	2.3	4.2
9 th Rib Fx	13	1.6	9 th Rib Fx	7	0.9	2.5
10 th Rib Fx	8	1.0	10 th Rib Fx	4	0.5	1.5
11 th Rib Fx	1	0.1	11 th Rib Fx	3	0.4	0.5
12 th Rib Fx	1	0.1	12 th Rib Fx	0	0	0.1

Table 1: Frequency of Radiologic Findings of the Studied Patients.

Mean mediastinal width of patients was 80.9±11.45 (49.2-142.29) mm with a median of 79.43 mm. Mediastinal width was not significantly different by gender ($P=0.36$), but was significantly associated with age ($P<0.001$).

Pearson chi square test showed the association of cause of trauma with pathologies as follows: right and left pneumothorax ($P=0.873$ and 0.023), right and left hemothorax ($P=0.115$ and 0.998), diaphragm and clavicle problems ($P=0.273$ and 0.677 , respectively), and rib fractures were not associated with cause of trauma, except right rib fracture ($P<0.001$). Right 9th and 2nd rib fracture was associated with sex ($P=0.058$ and 0.043 , respectively), but other pathologies were not significantly correlated with patients' sex. Among cases with left pneumothorax 4 cases had car-car accident, 7 had motor-car accident, 16 stab wound, 8 cases car turnover, 6 falling down, 8 car-patient accident, 3 motor-patient accident, and 1 bicycle trauma, and 1 gunshot.

Regarding co-occurrence of pathologies, 7 cases (0.88%) had both right and left pneumothorax, 18 cases (2.27%) had both right and left hemothorax. 2 cases had right pneumothorax and right clavicle fracture, 1 case had right pneumothorax and left clavicle fracture, 12 cases had left pneumothorax and left clavicle fracture, 3 cases had right hemothorax and right clavicle fracture, and 7 cases had left hemothorax and left clavicle fracture.

4 cases had both right pneumothorax and right diaphragmatic abnormalities and 20 cases had both right hemothorax and right diaphragmatic abnormalities cases and 1 patient had both left pneumothorax and right diaphragmatic abnormalities, 3 patients had both left hemothorax and right diaphragmatic abnormalities. Right diaphragmatic abnormalities co-occurred with right clavicle fracture in 3 patients and none cases of left clavicle fracture.

Left diaphragmatic abnormalities co-occurred with right hemothorax in 11 cases, right pneumothorax in 2 cases, left pneumothorax in 11 cases, left hemothorax in 1 case, right clavicle fracture in 4 and left clavicle fracture in 1 patient.

Discussion

As the results of the current study, consistent to other studies [21-23], have demonstrated, most traumatized patients are young men who are considered the productive group in the society and performing CXR routinely for all traumatized patients entering the emergency room exposes them to a high radiation rate and increases health costs. Therefore, we have performed CXR selectively for indicated patients.

Traumatic injuries has scarcely been studied in Iran. A recent study in Shiraz declared a mean age of 26.6 years including 75.8% men and have reported 22.9% car accident as the commonest cause of injury [21]. Although they have not excluded chest trauma, their results are comparable to the current study. Another study in Zahedan assessed 768 traumatic patients with a mean age of 23 including 80% men and have similarly reported road traffic accidents as the most frequent cause (about 60%) [22]. The studied patients in the two above-mentioned studies were apparently younger and they have not separated the forms of car accidents to be comparable with our results. A wide-scaled study has evaluated 58013 traumatic patients in Tehran (48173 blunt trauma) and has reported a mean age of 27 years with a male-to-female ratio of 4:1 and have reported the most common causes of injuries as gunshot, traffic accidents involving pedestrians, and falling in hospitalized patients [23], which was also in accordance with our results, as we also had a 4.78:1 male-to-female ratio. Although they have also not reported different statistics by body regions, they have reported

Citation: Mohammadian M, Shakibafard A, Akhavan M, Ghahramani Z, Paydar S (2017) Assessing Radiologic Findings of Chest Radiographs in Patients with Trauma Admitted to the Emergency Room. *Emerg Med Inves: EMIG-155*.

thorax as the third rank in most severe injuries. A population-based study in 2003 also reported road traffics in 1071 million males as the highest cause of disability premature deaths in Iran [24]. Parallel to the results of the current study, the most traumatized patients are young male who are considered the productive group in the society. Thus, paying attention to their initial care is necessary to provide the best approach for diagnosis and treatment and increase their survival.

Considering the radiologic findings, the commonest pathologies in the current study included rib fractures (42.7%), pneumothorax (11.6%), abnormal diaphragmatic findings (10.3%), and hemothorax (8.3%). Similarly have other studies reported different rates of abnormal CXR findings [9,17,25]. The rate of pneumothorax in the current study was lower than the mentioned studies, but as they also confirmed, pneumothorax can become a dangerous entity for the patient and its early diagnosis is essential in traumatic patients and studies have introduced different imaging methods as the most proper, accurate, and rapid imaging technique for its diagnosis; some have suggested CT [15], while others have concluded similar results reported by CXR and ultrasound [18]. As long as CT scan has a high cost, radiation rate, and long duration of imaging time and may be unavailable in all centers of our country, the trauma guideline of Rajaie hospital has considered CXR as the initial assessment of traumatic patients.

As studies have confirmed, selective CXR, as performed in the present study, reduces the health costs and radiation amount and is a safe and efficacious policy that prevents further investigations [8,10], compared to routinely performed CXR for all patients that has low clinical value [12], especially in stable conscious trauma patients with normal physical examination [13]. As long as there is no consensus on the gold standard imaging method, the trauma guideline of Rajaie hospital has considered selective CXR as the initial assessment of traumatic patients.

To the best of our knowledge, this study is the first study that evaluated the details about the cause of trauma in chest traumas and the comprehensive causes of road accidents and trauma in the current study help the decision-makers to have a better view.

The present study had some strengths including controlling errors of the study, for instance the emergency medicine specialist who read the radiographs initially in the emergency ward was blind to the case.

Beside the advantages of this study, it also faced some limitations, including not considering other socio-economic, psychologic, and other demographic details of the patients. As far as the study was performed in the emergency ward and there was limited time for gathering further details of the patient, as the patient should have been diagnosed and treated rapidly. It would also be greater, if we could compare the results of radiographies with CT scans and ultrasonography that were not performed as

they are not included in the trauma guideline of Rajaie hospital. We also did not have the choice to follow the patients, to observe if they required repeating the imaging or what further interventions they needed. Future studies are needed to give the emergency medicine specialists a better view in diagnosis and management of multiple traumatic patients.

The results of the current study, in accordance to previous studies, suggest rib fractures, pneumothorax, abnormal diaphragmatic findings, and hemothorax as the most frequent findings in CXR that need to be assessed meticulously. Moreover, the majority of patients were young males and the most frequent causes of trauma included car accidents, falling down, and stab wound. Therefore, paying attention to their diagnosis and treatment may increase the survival of this important group of patients.

References

1. Baker CC, Oppenheimer L, Stephens B, Lewis FR, Trunkey DD (1980) Epidemiology of trauma deaths. *The American Journal of Surgery* 140: 144-150.
2. Mehrdad R (2009) Health system in Iran. *JMAJ* 52: 69-73.
3. Zadeh H, Vahabi R, Nazparvar B, Amoei M (2002) An epidemiological study and determination of causes of traffic accident-related deaths in Tehran, Iran (during 2000–2001). *Journal of clinical forensic medicine* 9: 74-77.
4. LoCicero 3rd J, Mattox K (1989) Epidemiology of chest trauma. *The Surgical clinics of North America* 69: 15-19.
5. Collicott PE (1992) Advanced Trauma Life Support (Atls): Past, Present, Future-16th Stone Lecture, American Trauma Society. *Journal of Trauma and Acute Care Surgery* 33: 749-753.
6. Bouillon B, Kanz K, Lackner C, Mutschler W, Sturm J (2004) The importance of advanced trauma life support (ATLS) in the emergency room. *Der Unfallchirurg* 107: 844-850.
7. Wall B, Hart D (1997) Revised radiation doses for typical X-ray examinations. Report on a recent review of doses to patients from medical X-ray examinations in the UK by NRPB. National Radiological Protection Board. *The British journal of radiology* 70: 437-439.
8. Paydar S, Ahmadi A, Dalfardi B, Shakibafard A, Abbasi H, et al. (2014) Clinical and economic effects of selective radiological evaluation of high-energy trauma patients: a prospective experience of a level 1 busy trauma centre. *Emergency Medicine Journal* 204083.
9. Aukema TS, Beenen LF, Hietbrink F, Leenen LP (2012) Initial assessment of chest X-ray in thoracic trauma patients: Awareness of specific injuries. *World journal of radiology* 4: 48.
10. Sears BW, Luchette FA, Esposito TJ, Dickson EL, Grant M, et al. (2005) Old fashion clinical judgment in the era of protocols: is mandatory chest X-ray necessary in injured patients? *Journal of Trauma and Acute Care Surgery* 59: 324-332.
11. Wisbach GG, Sise MJ, Sack DI, Swanson SM, Sundquist SM, et al. (2007) What is the role of chest X-ray in the initial assessment of stable trauma patients? *Journal of Trauma and Acute Care Surgery* 62: 74-79.

Citation: Mohammadian M, Shakibafard A, Akhavan M, Ghahramani Z, Paydar S (2017) Assessing Radiologic Findings of Chest Radiographs in Patients with Trauma Admitted to the Emergency Room. *Emerg Med Inves: EMIG-155*.

12. Paydar S, Johari HG, Ghaffarpasand F, Shahidian D, Dehbozorgi A, et al. (2012) The role of routine chest radiography in initial evaluation of stable blunt trauma patients. *The American journal of emergency medicine* 30: 1-4.
13. Forouzanfar MM, Safari S, Niazazari M, Baratloo A, Hashemi B, et al. (2014) Clinical decision rule to prevent unnecessary chest X;ray in patients with blunt multiple traumas. *Emergency Medicine Australasia* 26: 561-566.
14. Rodriguez RM, Hendey GW, Marek G, Dery RA, Bjoring A (2006) A pilot study to derive clinical variables for selective chest radiography in blunt trauma patients. *Annals of emergency medicine* 47: 415-418.
15. Trupka A, Waydhas C, Hallfeldt K, Nast-Kolb D, Pfeifer K, et al. (1997) Value of thoracic computed tomography in the first assessment of severely injured patients with blunt chest trauma: results of a prospective study. *Journal of Trauma and Acute Care Surgery* 43: 405-412.
16. Exadaktylos AK, Sclabas G, Schmid SW, Schaller B, Zimmermann H (2001) Do we really need routine computed tomographic scanning in the primary evaluation of blunt chest trauma in patients with "normal" chest radiograph? *Journal of Trauma and Acute Care Surgery* 51: 1173-1176.
17. Traub M, Stevenson M, McEvoy S, Briggs G, Lo SK, et al. (2007) The use of chest computed tomography versus chest X-ray in patients with major blunt trauma. *Injury* 38: 43-47.
18. Ma OJ, Mateer JR (1997) Trauma ultrasound examination versus chest radiography in the detection of hemothorax. *Annals of emergency medicine* 29: 312-316.
19. Blaivas M, Lyon M, Duggal S (2005) A prospective comparison of supine chest radiography and bedside ultrasound for the diagnosis of traumatic pneumothorax. *Academic Emergency Medicine* 12: 844-849.
20. Shapiro M, Heiberg E, Durham R, Luchtefeld W, Mazuski J (1996) The unreliability of CT scans and initial chest radiographs in evaluating blunt trauma induced diaphragmatic rupture. *Clinical radiology* 51: 27-30.
21. Abbasi HR, Mousavi SM, Akerdi AT, Niakan MH, Bolandparvaz S, et al. (2013) Pattern of traumatic injuries and Injury Severity Score in a major trauma center in Shiraz, Southern Iran. *Bulletin of Emergency and Trauma* 1: 81-85.
22. Chardoli M, Rahimi-Movaghar V (2006) Analysis of trauma outcome at a university hospital in Zahedan, Iran using the TRISS method. *East African medical journal* 83: 440-442.
23. Zargar M, Modagheh M-HS, Rezaishiraz H (2001) Urban injuries in Tehran: demography of trauma patients and evaluation of trauma care. *Injury* 32: 613-617.
24. Naghavi M, Abolhassani F, Pourmalek F, Lakeh MM, Jafari N, et al. (2009) The burden of disease and injury in Iran 2003. *Population health metrics* 7: 9.
25. Ball CG, Ranson K, Dente CJ, Feliciano DV, Laupland KB, Dyer D, et al. (2009) Clinical predictors of occult pneumothoraces in severely injured blunt polytrauma patients: a prospective observational study. *Injury* 40: 44-47.