

Research Article

Situation of Sexually Transmitted Infections (STIs) in Sudan

Nasr Abdalla Mohamed Elawad*

Department of Obstetrics and Gynecology, Sudan Fertility Care Association, Sudan

*Corresponding author: Nasr Abdalla Mohamed Elawad, Department of Obstetrics and Gynecology, Sudan. Tel: +249912358430; Email: Nasr.abdalla@gmail.com

Citation: Elawad NAM (2018) Situation of Sexually Transmitted Infections (STIs) in Sudan. Adv Reprod Sci Reprod Health Infertil: ARRHI-102. DOI: 10.29011/ARRHI-102. 100002

Received Date: 28 February, 2018; Accepted Date: 21 March, 2018; Published Date: 30 March, 2018

Abstract

In general, STIs are epidemics and present an enormous health and economic consequences. An estimated half a billion new cases of STIs occur each year globally, with women being more susceptible and bearing the greatest burden of these infections. Sudan geographically is bounded by nine countries most of them have a high prevalence of HIV seropositive. Currently, the HIV seropositive prevalence in Sudan is within the range of 1.2-1.6% but is expected to increase as the present political and security instability in some of the neighbouring countries led to influx of refugees to Sudan. The refugees are now living among the low socioeconomic groups and represent a significant fragment of non-skilled labour forces.

STIs are a public health concern, but in Sudan very few studies have measured the prevalence of STIs nationwide. Based on the results of the conducted surveys, the populations' awareness on STIs is low and they are large differentials by education and residence. Sudan government in collaboration with UN agencies and NGOs is implementing certain interventions to prevent and control STIs including HIV/AIDS.

A program of syndromic management of STIs is implemented at the primary health care levels. The STIs situation needs more and great efforts from the stakeholders and the government should encourage and support studies on STIs. Based on that, an appropriate and evidence-based strategy can be developed to prevent and control STIs including HIV/AIDS.

Objective: The objective of this study is to assess the prevalence of and knowledge on STIs in Sudan.

Keywords: Chlamydia; Gonorrhea; Prevalence; STIS Other than HIV/AIDS; Syphilis; Sudan; Trichomoniasis

Methodology

Literature review was conducted through electronic searches using the words (STIs in Sudan). In addition; the national published documents with relevant data on STIs were also searched. The findings of this study are a result of a review of 22 documents.

General information

Sudan is a vast country in sub-Saharan Africa with a total population of around 39.2 million persons. Of the total population, rural population formed about 63.2%, urban 29.8% and nomads 7 % [1]. Sudan has a prominent diversity in, culture, religion, languages and ethnicity. Sudan has experienced long-term ethnic and political conflicts; some of them are still ongoing in Dar Fur,

South Kordofan and the Blue Nile states. Sudan is bounded by nine countries, some of them having a high prevalence of HIV infection and acquired AIDS. Due to political instability and conflicts in some of the Sudan neighbouring countries, Sudan currently hosts around two million refugees reside in the eastern and southern parts of Sudan, some of them engaged in the labour forces. This situation makes Sudan at risk for an increase in the prevalence of STIs including HIV/AIDS. The first case of HIV/AIDS in Sudan was reported in 1986 and in 2002 the total number of cases reported had increased to 4004, representing a seropositive prevalence of 1.6%. The main mode of HIV transmission is heterosexual accounting for 97% of HIV positive cases [2]. STIs are a major public health problem in all regions of the world. WHO estimated in 2005 that about 448 million curable infections occur every year worldwide in adult men and women [3].

Sexually Transmitted Infections (STIs) are defined as

infections that spread primarily through person-to-person sexual contact. However, other infections, in particular HIV/AIDS, syphilis and hepatitis B, can also be transmitted via mother-to-child-transmission during pregnancy, childbirth and lactation, blood products and tissue transfer. There are more than 30 different sexually transmitted bacteria, viruses and parasites responsible for STIs [4]. STIs should be distinguished from Reproductive Tract Infections (RTIs). RTIs are defined as infection of the genital organs and include endogenous infection such as bacterial vaginosis and vulvovaginitis candidiasis. These two infections are mostly not sexually transmitted, and they can occur in women who have never had a sexual relationship. Some factors may increase the risk of endogenous infections e.g. intake of antibiotics or contraceptive pills; pregnancy; uncontrolled diabetes or low immunity system [5].

STIs and their complications are among the most important causes of illness and death for women in the developing countries. STIs are associated with an increased risk of both acquisition and transmission of HIV. Untreated chlamydia infection is estimated to be the cause of at least a third of female infertility. Also, women who have had Pelvic Inflammatory Diseases (PIDs) are six to ten times more likely to have an ectopic pregnancy than those who have not had one. Untreated maternal syphilis infection may lead to stillbirth and neonatal deaths. Up to 35% of pregnancies among women with untreated gonococcal infection result in spontaneous abortions, premature deliveries, and up to 10% of perinatal deaths [4].

In some women, gonorrhea symptoms are so mild that they go unnoticed. Many women with gonorrhea discharge think they have a yeast infection and self-treat with medications purchased over the counter. Because vaginal discharge can be a sign of a number of different problems, it is best to always seek the advice of a doctor to ensure correct diagnosis and treatment [6]. Gonorrhoea may also be spread by contact with infected bodily fluid, so that an infected mother can pass on the infection to her newborn during childbirth. Once the gonorrhea bacteria come into contact with the eyes of the newborn, the process may end with acute conjunctivitis [7].

In addition, many people are infected with non-curable STIs, mainly viral diseases such as HIV/AIDS, hepatitis B or genital herpes. About 536 million people aged 15-49 were estimated to be living with herpes simplex virus type 2 worldwide in 200 [8].

Knowledge and Modes of Transmission of STIs in Sudan

Information available on STIs in Sudan is very limited. The social stigma attached to these diseases, prevents proper dissemination of basic knowledge about the disease. Information was sought on knowledge of ever-married women on HIV/AIDS, and two symptoms of other STIs [5]. The survey sought to know whether women have heard of these conditions, and their knowledge about mode of transmission, preventive and curative measures.

Overall, 43% of women have heard of AIDS. The awareness of AIDS is particularly poor among rural women, only 28% of them have heard of AIDS. There are also large differentials in knowledge by education, 95% of women with secondary education and above have heard about AIDS compared with 22% of illiterate women [9].

91% of women who have heard about AIDS reported sexual intercourse as the main mode of transmission. An overwhelming majority 83% has reported that AIDS could be avoided by having only one sexual partner. 94% suggested use of disposable syringes, and 37% use of condoms as methods of STIs prevention [9]. Only 30% of women heard about genital ulcer. With regard to mode of HIV transmission, 59% of women reported sexual intercourse as a main mode [5]. Quite a few women had misconceptions about the mode of HIV transmission, such as contaminated toilet (24%), contact with AIDS patients (17%). 51% of women reported awareness of abnormal vaginal discharge. 42% of rural women are aware about abnormal vaginal discharge compared with 66% of urban women [9].

A survey conducted in 2006 shows 70% of women age 15-49 years heard about AIDS, awareness about AIDS is low among illiterate women compared with educated ones, 49.6% vs 90.5 % [6]. Regarding mode of HIV transmission 51.5% admitted sexual intercourse, 39.7 through contaminated blood transfusion and 38.8 through injectables. Only 7.5% of women mentioned use of condoms protects HIV transmission, 12.9% mentioned insect bite can transmit HIV and 1.6% stated that sharing foods and eating with people affected by AIDS can help HIV transmission. 54% of women are aware about HIV vertical transmission. Awareness of vertical HIV transmission is 29.7% among illiterate women vs. 85% among literate ones [10].

87% of men age 15-24 years has heard of HIV/AIDS which is similar to men age 15-49 years, and the comprehensive knowledge about HIV prevention is 11% for men age 12-24 and 12% of those age 15-49 years [11]. A cross-sectional study on knowledge of AIDS was carried out in 2007 among a census of dental students in six dental faculties in Khartoum, the capital of Sudan. Lectures and Radio/TV were the most frequently reported sources of information related to HIV/AIDS as reported by 61% and 44% of the students. 47.6% of students confirmed a need for further education across topics related to HIV/AIDS. 97% of students mentioned HIV transmission through contaminated blood transfusion. About half the students recognized condoms use as a safe method of preventing transmission of HIV [12].

The Epidemiology of STIs in Sudan

Although STIs cause much morbidity, mental ill-health and social problems, yet they are currently being neglected in many developing countries. There are few facilities for correct diagnosis and treatment. In Sudan case, there are no adequate statistics and

enough studies, have been done to determine the dimensions of STIs status. To estimate the prevalence of STIs among women in a Sudanese community, 338 women with ages ranging from 15 to 69 years were randomly selected and studied. The results showed that trichomoniasis was found in 7.7%, gonorrhoea in 1.2%, HIV in 1.2% and syphilis in 0.9% of the subjects [13].

A study was conducted in two venereal diseases clinics in Khartoum (the capital of Sudan) to assess the STIs in Sudanese males. Out of the total 138 patients referred to the clinic with a referral diagnosis of STIs, 61 patients (47.1%) were found not to be suffering from any STI.

Among male adult patients, Non-Gonococcal Urethritis (NGU) was the commonest STI encountered (35.1%), next came gonococcal urethritis (25.9%). Syphilis accounted for only 1.3% of the cases investigated. Most of the patients with STIs were in the age range 20-39 years. Of the infected patients 49.3% had their infections from prostitutes. Nearly half of the patients examined and found infected with STIs were in the low-paid socioeconomic group; and 71.4% of them were single [14].

Underprivileged pregnant women attending antenatal clinic for routine checkups in displaced camps, a women's prison and several peripheral health centres were clinically and laboratory screened for STIs other than HIV/AIDS. A total of 426 women with an age range of 14-45 were included. Clinical data, blood, cervical and vaginal swabs were collected. All attendees were HIV/1/2-negative. The prevalence of *Trichomonas vaginalis* was found to be 7.8%, *Chlamydia trichomatis* 49%, *Neisseria gonorrhoea* 0% and *Treponema pallidum* 5%. Although vaginal discharge, among other symptoms, is known to be the most significant indicator for STIs, in this study the authors stated that their identified predictive value was only 14.1%. They concluded that the use of syndromic approach for diagnosing and treating attendees of antenatal settings is of low clinical value and many easily curable STIs will be overlooked [15].

The number of HIV infected cases is reported to have risen to more than 7,245 in 2000 of which there are 3,638 AIDS cases [9]. In all these cases, heterosexual activity accounts for more than 95.5%. The possibility of further increase of STIs in Sudan is higher as Sudan shares border with nine African countries, 6 of which have the highest prevalence of STIs/ HIV/AIDS.

STIs Prevention and Control

The government of Sudan includes STIs in RH top priorities since 1990s. Currently due to unexplained reasons the government is focusing mainly on HIV/AIDS issues and to some extent hepatitis B virus and syphilis. The Sudan National AIDS Program (SNAP) is the technical body with the responsibility for national policy, planning and coordination.

With a focus on reducing HIV transmission and HIV mortality, and under SNAP control and with support of the United Nations Development Program (UNDP), by the end of 2011, there were 144 VCT centres; 30 ART centres and 80 PMTCT sites [16].

The services in all these sites are free and based on WHO guidelines and recommendations. Considering the relative advantage of reaching the community, since 1996 the NGOs in Sudan are involved in the HIV activities, particularly raising awareness and conducted selected outreach interventions for HIV key and vulnerable populations. In spite of these efforts, recent research suggested that HIV/AIDS continues to spread in Sudan with a prevalence likely to reach 1.2 per cent of the population by 2015, almost double what it was in 2009 [17]. Other than HIV/AIDS, the most widely known STIs are gonorrhoea, syphilis, chlamydia and trichomoniasis. The relation between HIV and other STIs makes it even more urgent to prevent and control curable STIs. Syndromic case management is consistently applied at any first-level health facility, such as health centre, rural hospital or STIs clinic.

The result of integrating the management of STIs in the primary health care and adoption of syndromic case management, the number of STIs treated cases increased from 35,263 in 2009 to 89,625 in 2011 [16].

Results

In general, STIs are epidemics and present an enormous health and economic consequences. Currently the prevalence of HIV infection in Sudan is within a range of 1.2-1.6% and expected to increase. The main mode of HIV transmission is heterosexual accounting for 97% of HIV positive cases. The awareness on AIDS is poor especially among women in rural areas. The prevalence of trichomoniasis was found in 7.7%, gonorrhoea in 1.2%, HIV in 1.2% syphilis in 0.9%, non-gonococcal urethritis 35.1%, and *Chlamydia trichomatis* 49%. Syndromic case management is consistently applied at all first-level health facilities.

Conclusion

Sexually Transmitted Infections (STIs) are important public health problem and major health threats affecting people globally; however, the burden of STIs is greatest in low-income countries including Sudan. There are few data on STIs other than HIV/AIDS in Sudan. The current prevalence of HIV/AIDS is low but, expected to increase. The most accounted STIs among the studied groups are gonorrhoea, trichomoniasis, chlamydia, syphilis and HIV/AIDS. The most frequent transmission mechanism of STIs, in Sudan, is heterosexual relations. Syndromic approach is commonly used for diagnosing and treating STIs other than HIV/AIDS.

References

1. Sudan Census (2008).

2. UN AIDS U, WHO, (2004) Assessment of the epidemiological situation, UNAIDS.
3. WHO (2011) Prevalence and incidence of selected sexually transmitted infections. Geneva: WHO.
4. WHO (2013) Report on global STIs surveillance.
5. www.fairview.org/Healthlibrary/article/116682AA.
6. Sexual conditions guide.
7. www.fpa.org.uk/sites/default/files/gonorrhoea.
8. An estimate of the global prevalence and incidence of herpes simplex virus type 2 infections (2008) *Bull WHO* 86: 805-812.
9. Sudan Safe Motherhood Survey (SMS) (1999).
10. Sudan Household Survey (SHHS) (2006).
11. Sudan household Survey (SHHS) (2010).
12. Elwalid F. Nasir, Anne N. Astorm, Jamil David and Rouf W Ali (2008) HIV and AIDS related knowledge, sources of information, and reported need for further education among dental students in Sudan- a cross sectional study. *BMC Public Health* 8: 286.
13. Kafi SK, Mohamed AO, Musa HA (2000) Prevalence of sexually transmitted diseases (STD) among women in a suburban Sudanese community. *Ups J Med Sci* 105: 249-253.
14. Omer EE, Ali MH, Taha OM, Ahmed MA, Abbaro SA (1982) Sexually transmitted diseases in Sudanese males *Trop Doct* 12(4 Pt 2): 208-210.
15. *Int J STD Online* first (2011).
16. Sexually transmitted infections other than HIV/AIDS among women of low socio-economic class attending antenatal clinics in Khartoum, Sudan (2016).
17. Fighting HIV/AIDS in Sudan.