Growth Hormone-Releasing Hormone (GHRH): Its Biology and Therapeutic Aspects

Liting Zheng*

College of Animal Science and Veterinary Medicine, Shenyang Agricultural University, Shenyang, Liaoning, China

*Corresponding author: Liting Zheng, College of Animal Science and Veterinary Medicine, Shenyang Agricultural University, Shenyang, Liaoning 110011, China. Email: liting_zheng@yahoo.com

Citation: Zheng L (2018) Growth Hormone-Releasing Hormone (GHRH): Its Biology and Therapeutic Aspects. J Pharma Pharma Sci: JPPS-174. DOI: 10.29011/2574-7711.100074

Received Date: 01 October, 2018; **Accepted Date:** 15 August, 2018; **Published Date:** 23 August, 2018

Introduction

GHRH, (also known somatocrinin; Growth Hormone Releasing Factor (GRF); Growth hormone releasing hormone (GRH) and somatostatin) is a 44-amino acid peptide which synthesized in arcuate nucleus of the hypothalamus in Central Nervous System (CNS) and subsequently released into hypothalamic-pituitary portal vascular system. By binding with GHRH receptors at anterior pituitary gland, GHRH stimulates synthesis and release of Growth Hormone (GH). After Schally, AV et al made his discoveries concerning the peptide hormone production of the brain [1-3], in 1982, GHRH was isolated and sequenced its full length by two groups Rivier, J et al and Guillemin, R et al. from two pancreatic tumors patients beared with acromegaly [4,5]. All these discoveries made GHRH knowledge more rapidly acquired by scientists in the following decades. The important roles GHRH played in different biological process, such as energetic metabolism [6], cancer initiation and progression [7], sleep process [8], pituitary hypoplasia and dwarfism [9], also gradually recognized by researchers.

Mayo KE. et al. mapped the first GHRH gene to chromosome 20 and reported it is synthesized as 108 amino-acid preproGHRH. After processed by unknown proteinases in two cleavage points (Arg-Arg and Gly-Arg in Figure 1), human pituitary GHRH1-44 was then released, whose biological activity is almost the same as its C-terminal truncated derivatives, which are due to alternative processing in its preproproteins [10,11]. Further studies revealed that removal of C terminal amino acids and its N-terminal

molecules from GHRH1-29 will both result in dramatic loss of its GH releasing activity and less rapid degradation rate, suggesting the importance of these amino acids for peptide structure and stability [11]. Up to now, no mutations have been reported regarding the GHRH genes.

```

MPLWVFFFVILTLNSSHCSPPPPLTRMRRYADAIFTNSYR
      |                               |
      Signal peptide                pro peptide
KVLGQLSARKLLQDIMSRQQGESNQERGARARLGRQVDS
      |                               |
      GHRH1-29                      GHRH1-44
MWAEQKQMELESILVALLQKHSRNSQG

```

Figure 1: Amino acids sequence of human preproGHRH. Red letters represent possible cleavage sites.

Expression of GHRH has been found in various normal extra-pituitary tissues: ovary, placenta, testis, pancreas, gastrointestinal tract, prostate and immune cells (Figure 2) and biological fluids, such as CSF (human cerebrospinal fluids) and milk [12]. The content of GHRH in the human hypothalamus is vary from 10-250ng/hypothalamus and has a low level in peripheral plasma, but when ectopic GHRH-secreting tumors occurred; the content in plasma will be increased indicating a promising biomarker for diagnosis of specific tumors. The structure of GHRH from several species exhibits almost complete homology except the rats. The GHRH in rats only has 70% homology with humans and 43-amino acids in its full biological activity form. The difference mainly resides in C terminal sequence, which results in decreased biological activity and receptor binding ability.

Figure 2: GHRH gene expression of different normal human tissues. (analysis from gene card) Left panel: the results from RNA seq analysis. Middle panel: the results from Microarray analysis. Right panel: the predict results from SAGE (serial analysis of gene expression) analysis.

GHRH exerts its function by binding to GHRH receptors (GHRHRs), which is belonged to the Gs-protein – couple receptors. After binding to GHRHRs, GHRH stimulates the Adenylyl Cyclase (AC) with resulting increase in cAMP levels and activation of proteinase A. Then, with the influx of calcium, GH is released from somatotrophs. It is also reported that by binding with GHRHRs, the down streaming signals are associated with MAPK pathways, STAT3-PAK1 pathways and cell cycle factors [7].

GHRH Therapeutic Aspects

Physiologically, GH deficiency is due to familial GH deficiency (5%-10%), autosomal recessive mutations; and impaired GH response to GHRH. Clinically, the patients with irreversible GH response to GHRH include patients bear with organic pituitary disease, thalassemia or sideroblastic anemia [13] and primary microcephaly [14]. The reason for this is partly due to a decreased somatotroph secretory reserve, decreased pituitary GH stores or a resistance to the action of GHRH, the decreased content iron deposition in the pituitary and mutated PLEKHG2 with impaired actin polymerization. However, the patients with hypothalamic disease and hypothyroidism represent a reversible response of GH

to GHRH, and a recovery of this response will always be seen after initiation of treatment with thyroid hormone in patients [13]. Sermorelin, a functional peptide fragment of GHRH, has been used in the diagnosis of deficiencies in growth hormone secretion and measuring GHRH plasma levels therefore provides a precise and cost-effective test for the diagnosis of ectopic acromegaly. At present, it has been clearly demonstrated that GHRH and/or GHS represent reliable tools for the diagnosis of GHD [15,16].

Obesity patients showed a blunted GH response to all CNS-mediated stimuli, and it will be significantly improved by weight loss. For the diabetes patients, agonists of Growth Hormone-Releasing Hormone (GHRH) have been previously reported to promote growth, function, and engraftment of islet cells following transplantation, in 2013, Zhang et al reported that transplantation of rat islets preconditioned in vitro with GHRH agonists and its administration in vivo promoted growth, function, and engraftment of exogenous islets [6], supporting the use of GHRH agonists in type-1 diabetes.

The discovery of local autocrine/paracrine production of GHRH and other cells, directed the research to synthesis of more

potent GHRH antagonists to strongly inhibit the tumor growth with scarce endocrine action. The receptor mediated mechanisms comprise complex and still not completely understood on intracellular pathways that are strictly related to human tumorigenesis. However, it creates a new potential therapeutic option for various neoplasms [7]. The growing knowledge of mechanisms gives us new ways to its possible applications. Tesamorelin, a GHRH analog, trade name Egrifta, received U.S. Food and Drug Administration approval in 2010 for the treatment of lipodystrophy in HIV patients under highly active antiretroviral therapy [17].

In summary, new understandings have been gained in relation to the pathogenesis with different diseases of GH deficiency and GH excess state, and the use of GHRH and its analogs as diagnostic and therapeutic agents already represents a reality.

References

- Schally AV, Müller EE, Arimura A, Bowers CY, Saito T, et al. (1967) Releasing factors in human hypothalamic and neurohypophyseal extracts. *J Clin Endocrinol Metab* 27: 755-762.
- Schally AV, Bowers CY (1964) Corticotropin-Releasing Factor and Other Hypothalamic Peptides. *Metabolism* 13: SUPPL: 1190-1205.
- Schally AV, Kuroshima A, Ishida Y, Arimura A, Saito T, et al. (1966) Purification of growth hormone-releasing factor from beef hypothalamus. *Proc Soc Exp Biol Med* 122: 821-823.
- Rivier J, Spiess J, Thorner M, Vale W (1982) Characterization of a growth hormone-releasing factor from a human pancreatic islet tumour. *Nature* 300: 276-278.
- Guillemin R, Brazeau P, Böhlen P, Esch F, Ling N, et al. (1982) Growth hormone-releasing factor from a human pancreatic tumor that caused acromegaly. *Science* 218: 585-587.
- Zhang X, Cui T, He J, Wang H, Cai R, et al. (2015) Beneficial effects of growth hormone-releasing hormone agonists on rat INS-1 cells and on streptozotocin-induced NOD/SCID mice. *Proc Natl Acad Sci U S A* 112: 13651-13656.
- Wang H, Zhang X, Vidaurre I, Cai R, Sha W, et al. (2018) Inhibition of experimental small-cell and non-small-cell lung cancers by novel antagonists of growth hormone-releasing hormone. *Int J Cancer* 142: 2394-2404.
- Obal F Jr, Krueger JM (2004) GHRH and sleep. *Sleep Med Rev* 8: 367-377.
- Sundralingam T, Tennekoon KH, de Silva S, De Silva S, Hewage AS (2017) Pathogenic and likely pathogenic genetic alterations and polymorphisms in growth hormone gene (GH1) and growth hormone releasing hormone receptor gene (GHRHR) in a cohort of isolated growth hormone deficient (IGHD) children in Sri Lanka. *Growth Horm IGF Res* 36: 22-29.
- Mayo KE, Vale W, Rivier J, Rosenfeld MG, Evans RM (1983) Expression-cloning and sequence of a cDNA encoding human growth hormone-releasing factor. *Nature* 306: 86-88.
- Siejka A, Lawnicka H, Melen-Mucha G, Motylewska E, Komorowski J, et al. (2012) Antineoplastic action of growth hormone-releasing hormone (GHRH) antagonists. *Recent Pat Anticancer Drug Discov* 7: 56-63.
- Zheng G, Wang H, Wei C, Li Y (2011) iGepros: an integrated gene and protein annotation server for biological nature exploration. *BMC Bioinformatics* 12 Suppl 14: S6.
- Frohman LA, Jansson JO (1986) Growth hormone-releasing hormone. *Endocr Rev* 7: 223-253.
- Edvardson S, Wang H, Dor T, Atawneh O, Yaacov B, et al. (2016) Microcephaly-dystonia due to mutated PLEKHG2 with impaired actin polymerization. *Neurogenetics* 17: 25-30.
- Aimaretti G, Baldelli R, Corneli G, Bellone S, Rovere S, et al. (2004) GHRH and GH secretagogues: clinical perspectives and safety. *Pediatr Endocrinol Rev* 2 Suppl 1: 86-92.
- Aguar-Oliveira MH, Davalos C, Campos VC, Oliveira Neto LA, Marinho CG, et al. (2018) Hypothalamic abnormalities: Growth failure due to defects of the GHRH receptor. *Growth Horm IGF Res* 38: 14-18.
- Mangili A, Falutz J, Mamputu JC, Stepanians M, Hayward B (2015) Predictors of Treatment Response to Tesamorelin, a Growth Hormone-Releasing Factor Analog, in HIV-Infected Patients with Excess Abdominal Fat. *PLoS One* 10: e0140358.